

Scheda tassonomica:	
Classis	Gastropoda
Ordo	Neogastropoda
Familia	Costellariidae
Subfamilia	
Genus	Vexillum
Subgenus	(Pusia)
specie	ebenus (Lamarck, 1811) [Mitra]
subspecie	

Scheda descrittiva:	
<b>Sinonimi</b>	! <i>Voluta caffra</i> Linné, 1758 -- - sensu Olivi, 1792
	! <i>Voluta vulpecula</i> - Linné, 1758 -- - sensu Renier, 1804
	= <i>Mitra plumbea</i> - Lamarck, 1811
	= <i>Voluta plicatula</i> - Brocchi, 1814
	= <i>Mitra biplicata</i> - Risso, 1826 -: p. 242 (original description)/type locality: Alpes-Maritimes, France
	= <i>Mitra cornicula</i> - Risso, 1826 [Has homonym[s]]
	= <i>Mitra costulata</i> - Risso, 1826 -: p. 246, pl. 5 fig. 66 (original description/type locality: Alpes-Maritimes, France
	= <i>Mitra defrancei</i> - Payraudeau, 1826
	= <i>Mitra littoralis</i> - Risso, 1826
	<i>Mitra littoralis</i> - Risso, 1826 -- - Arnaud, 1978 : (lectotype designated)
	! <i>Mitra cornicularis</i> - Lamarck, 1822 -- - sensu Costa O.G., 1829
	<i>Mitra cordierii</i> - Maravigna, 1853 -: p. 137 (original description)/type locality: not specified [Sicily]
	= <i>Mitra ebenus</i> var. <i>costata</i> - Weinkauff, 1868
	= <i>Mitra ebenus</i> var. <i>laevigata</i> - Weinkauff, 1868
	<i>Mitra ebenus</i> var. <i>bifasciata</i> - Monterosato, 1875 -: (nomen nudum)
	<i>Mitra ebenus</i> var. <i>concolor</i> - Monterosato, 1875 -: (nomen nudum)
	<i>Mitra ebenus</i> var. <i>lanceolata</i> - Monterosato, 1875 -: (nomen nudum)
	<i>Mitra ebenus</i> var. <i>oblonga</i> - Monterosato, 1875 -: (nomen nudum)
	<i>Mitra ebenus</i> var. <i>turrita</i> - Monterosato, 1875 -: (nomen nudum)
	= <i>Mitra congesta</i> - Locard, 1886 -: p. 105-106, 540-541 (original description)/type locality: not specified [Mediterranean coast of France]
	= <i>Mitra ebenina</i> - Locard, 1886 -: p. 104 (unjustified emendation suggested in synonymy; made available by usage as valid name in Locard, 1892: 47)
	! <i>Voluta pyramidella</i> - Brocchi, 1814 -- - sensu Locard, 1886 : p. 105
	= <i>Mitra gracilis</i> - Locard, 1890 -: p. 109 (original description)/type locality: Mediterranean coast of France
	= <i>Mitra servaini</i> - Locard, 1890 -: p. 109 (original description)/type locality: Mediterranean coast of France
	= <i>Mitra bourguignati</i> - Locard, 1892 -: p. 48 (original description)/type locality: not specified [Mediterranean France]
	= <i>Mitra plicatuliformis</i> - Locard, 1892 -: p. 49 (original description)/type locality: not specified [French Mediterranean]
	= <i>Mitra subpyramidella</i> - Locard, 1892 -: p. 48-49 (original description)/type locality: not specified [French Mediterranean]
	= <i>Turricula</i> ( <i>Uromitra</i> ) <i>ebenus</i> var. <i>nigra</i> - Pallary, 1900 -: p. 264
	= <i>Mitra tunetana</i> - Pallary, 1903
	= <i>Uromitra ebenus</i> var. <i>totanigra</i> - Pallary, 1912
	? <i>Uromitra blanci</i> - Pallary, 1912 -: p. 20, pl. 1 fig. 54-55 (original description)/type locality: Tunis lagoon, Tunisia
	= <i>Mitra ebenus</i> var. <i>cesarea</i> - Parenzan, 1970 -: p. 191, pl. 40 fig. 765 [stated as new variety although published as a trinomen; hence not an available name]
	= <i>Mitra ebenus haifensis</i> - Nordsieck, 1982
<b>Descrizione</b>	La conchiglia è molto allungata, lucida. la columella è percorsa da 3 o 4 pieghe. La colorazione è marrone, con una linea chiara che percorre ogni giro. Il mollusco è carnivoro e vive nella sabbia. Comune.
<b>Locus typicus</b>	
<b>Descrizione originale</b>	
<b>Riferimenti bibliografici</b>	<a href="http://www.naturamediterraneo.com/forum/topic.asp?TOPIC_ID=4268">http://www.naturamediterraneo.com/forum/topic.asp?TOPIC_ID=4268</a> <a href="http://www.somali.asso.fr/clemam/biotaxis.php">http://www.somali.asso.fr/clemam/biotaxis.php</a>

Scheda fotografica:

Foto



f./var.  
provenienza  
misure  
Note

Foto di Aldo Marinelli